

BOSONIC FORMULAS FOR (k, l) -ADMISSIBLE PARTITIONS

B. FEIGIN, M. JIMBO, S. LOKTEV, T. MIWA AND E. MUKHIN

ABSTRACT. Bosonic formulas for generating series of partitions with certain restrictions are obtained by solving a set of linear matrix q -difference equations. Some particular cases are related to combinatorial problems arising from solvable lattice models, representation theory and conformal field theory.

1. INTRODUCTION

Let k, l, n be positive integers. A partition of length n

$$\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n \geq 0) = (0^{x_0}, 1^{x_1}, 2^{x_2}, \dots)$$

is called (k, l) -admissible if

$$\lambda_i \geq \lambda_{i+k} + l \quad (i = 1, \dots, n - k).$$

Then a partition λ is (k, l) -admissible if and only if the set of integers $(x_0, x_1, \dots)$ satisfies

$$0 \leq x_i \leq k, \quad x_i + \cdots + x_{i+l-1} \leq k. \quad (1.1)$$

for all $i \geq 0$.

The sequences satisfying (1.1) have been discussed in the literature. For example, the case $(k, l) = (1, 2)$ is related to the Rogers-Ramanujan identities and is used in the computation of 1-point correlation functions of the hard hexagon model [B]. The case where k is general and $l = 2$ appeared in the representation theory of the affine Lie algebra $\widehat{sl}_2$ in [FS]. More recently, the case $k = 2$ and $l \not\equiv 1 \pmod{3}$ appeared in the study of vertex operator algebras related to the $(2, 1)$ primary field in conformal field theory [FJM].

In this paper, we will not discuss such applications but treat the problem purely combinatorially for general k and l .

Given a positive integer N , consider (k, l) -admissible partitions of length n with $\lambda_1 < N$. Let $d_{k,l}^{(N)}(a, n)$ be the number of such partitions of weight a . In other words, $d_{k,l}^{(N)}(a, n)$ is the number of sequences $(x_0, \dots, x_{N-1}, 0, \dots)$, satisfying (1.1) such that

$$\sum_j jx_j = a, \quad \sum_j x_j = n.$$

In this paper we study the generating series of (k, l) -admissible partitions:

$$\chi_{k,l}(q, z) = \lim_{N \rightarrow \infty} \chi_{k,l}^{(N)}(q, z), \quad \chi_{k,l}^{(N)}(q, z) = \sum_{a,n} d_{k,l}^{(N)}(a, n) q^a z^n. \quad (1.2)$$

There are two different types of formulas which express the generating series of the kind (1.2) (one dimensional configuration sums): fermionic and bosonic formulas (see, e.g., [BMS]). In this paper, we present a bosonic formula in the infinite volume limit $N \rightarrow \infty$. The result is expressed as a finite sum of $(l-1)$ -fold series. Each series is a sum of terms parameterized by a set of $(l-1)$ non-negative integers; each term has the form

$$\frac{(-1)^\alpha q^\beta z^\gamma}{(q)_{t_2} \cdots (q)_{t_l} (q^{t_2 + \cdots + t_l} z)_\infty},$$

where $\alpha, \beta, \gamma, t_i$ are some integers described by explicit formulas.

It is worth mentioning the difference between our approach and that used, e.g., in [BMS]. Both approaches use finite polynomials $\chi_{k,l}^{(N)}$ and recurrence relations which connect N -th and $(N+1)$ -st polynomials. In [BMS] this recurrence is obtained by adding x_N to the sequences $(x_0, \dots, x_{N-1})$ with fixed boundary conditions at the *right end*, whereas in our approach, the recurrence is obtained by adding x_0 to sequences $(x_1, \dots, x_N)$ with fixed conditions at the *left end*.

In the latter approach of the right end recurrence, one can handle the generating series in a single variable q by setting $z = 1$. In the former approach of the left end recurrence, we have the shift of z to qz due to the shift of weight. The advantage of the left end recurrence is that it makes sense in the limit $N \rightarrow \infty$, where it becomes a q -difference matrix equation.

To solve the q -difference equation we use the following method. We consider the sum (1.2) as a sum over the polytope bounded by the conditions (1.1). In such a problem a classical idea is to express the result as a sum of contributions from the extremal points of the polytope. In general, it is not easy to enumerate all the extremal points of a polytope in large dimensions and compute the corresponding contributions. However, in our problem, the q -difference equation reduces this problem to the one-dimensional case: we construct the extremal points and the corresponding contributions for the $(N+1)$ -st polytope from those for the N -th.

The number of extremal points thus constructed grows exponentially in N . Miraculously, most terms cancel in pairs, and the final formulas contain only polynomially many terms. We do not understand the reason why this cancellation takes place. Because of technical restrictions, in this paper we give the result only in the limit $N \rightarrow \infty$.

A similar result by a similar method for the $\widehat{sl}_2$ coinvariants is obtained in [JMM].

The plan of the paper is as follows. In Section 2 we derive the q -difference equations and define operators A, B corresponding to creation of new extremal points. In Section 3 we determine the cancellation pairs. The proof of the cancellation is given in Section 4. Section 5 is devoted to describing the remaining terms after the cancellation.

2. THE (k, l) -CONFIGURATIONS AND THEIR CHARACTERS

2.1. The (k, l) -configurations. Let k, l be positive integers. An infinite sequence of non-negative integers $\mathbf{x} = (x_0, x_1, \dots)$ with finitely many non-zero entries is called a (k, l) -configuration if for all $j \in \mathbb{Z}_{\geq 0}$ we have

$$x_j + \dots + x_{j+l-1} \leq k. \quad (2.1)$$

The smallest non-negative integer N with the property $x_i = 0$ for all $i \geq N$ is called the *length* of the (k, l) -configuration $\mathbf{x}$.

Let $\mathbf{b} = (b_0, \dots, b_{l-2})$ be a vector with non-negative integer entries satisfying

$$b_0 \leq b_1 \leq \dots \leq b_{l-2} \leq k. \quad (2.2)$$

Denote by $\mathcal{C}_{k,l;\mathbf{b}}^{(N)}$ the set of all (k, l) -configurations of length at most N such that

$$x_0 \leq b_0, \quad x_0 + x_1 \leq b_1, \quad \dots, \quad x_0 + \dots + x_{l-2} \leq b_{l-2}.$$

We introduce the generating functions

$$\chi_{k,l;\mathbf{b}}^{(N)}(q, z) = \sum_{\mathbf{x} \in \mathcal{C}_{k,l;\mathbf{b}}^{(N)}} q^{\sum_{j \geq 0} j x_j} z^{\sum_{j \geq 0} x_j}, \quad (2.3)$$

and call them the *characters of (k, l) -configurations*.

Setting $i = x_0$ and considering a shifted configuration $x'_j = x_{j+1}$, we are led to the following recursion equations.

Lemma 2.1. *The set of characters $\chi_{k,l;\mathbf{b}}^{(N)}(q, z)$ ($N = 0, 1, \dots$) is the unique solution of the recursion equations*

$$\chi_{k,l;(b_0, \dots, b_{l-2})}^{(N+1)}(q, z) = \sum_{i=0}^{b_0} z^i \chi_{k,l;(b_1-i, b_2-i, \dots, b_{l-2}-i, k-i)}^{(N)}(q, qz), \quad (2.4)$$

with the initial condition $\chi_{k,l;\mathbf{b}}^{(0)}(q, z) = 1$ for all $\mathbf{b}$.

2.2. The operators A, B . Our main interest is to find a formula for the characters in “bosonic form”. To illustrate what it means, let us consider N points $\{0, 1, \dots, N-1\}$ on a line segment $[0, N-1]$. Define its character to be the sum $1 + z + \dots + z^{N-1}$. This polynomial can be written as a sum of two formal power series,

$$1 + z + \dots + z^{N-1} = \frac{1}{1-z} + \frac{z^{N-1}}{1-z^{-1}}, \quad (2.5)$$

where the right hand side is understood as an expansion in non-negative powers of z . The two terms correspond to the two end points (“extremal points”) of the segment. The contribution from each extremal point may be viewed as the character of a half line (or a cone) $\{i \geq 0\}$, $\{i \leq N-1\}$, respectively. We apply this principle to take the sum in the recursion equations (2.4).

Let P_i ($i = 1, \dots, l$) be monomials of q, z of the form $q^{m_i} z^{n_i}$ where $m_i, n_i \in \mathbb{Z}_{\geq 0}$. Introduce the notation

$$[P_1, \dots, P_l] = P_1^{b_0} P_2^{b_1-b_0} \dots P_{l-1}^{b_{l-2}-b_{l-3}} P_l^{k-b_{l-2}}.$$

We regard $[P_1, \dots, P_l]$ as a function of $q, z, \mathbf{b}, k$. Note that

$$[\lambda P_1, \dots, \lambda P_l] = \lambda^k [P_1, \dots, P_l].$$

Let V_l be the complex vector space spanned by functions $\xi(q, z; \mathbf{b}, k)$ of the form $f(q, z)[P_1, \dots, P_l]$, where $f(q, z)$ is a formal power series in q, z independent of $\mathbf{b}, k$, and $[P_1, \dots, P_l]$ belongs to the following list:

$$\begin{aligned} & q^{mk} z^{nk} [1, q^{c_1}, q^{c_2}, \dots, q^{c_{l-1}}], \\ & q^{mk} z^{nk} [q^{c_{l-1}} z, 1, q^{c_1}, \dots, q^{c_{l-2}}], \\ & \dots \\ & q^{mk} z^{nk} [q^{c_1} z, q^{c_2} z, \dots, q^{c_{l-1}} z, 1], \end{aligned}$$

with $c_i, m, n \in \mathbb{Z}_{\geq 0}$, $0 \leq c_1 \leq \dots \leq c_{l-1}$. We call such a function $\xi(q, z; \mathbf{b}, k)$ a simple vector, $[P_1, \dots, P_l]$ the *vector part* and $f(q, z)$ the *scalar part* of it.

Let S be the q -shift operator in z , defined by

$$(Sg)(q, z) = g(q, qz). \quad (2.6)$$

Define two linear operators A, B on V_l by the formulas

$$A(f(q, z)[P_1, \dots, P_l]) = S \left(\frac{f(q, z)}{1 - q^{-1} z P_l / P_1} [P_1, P_1, P_2, \dots, P_{l-1}] \right), \quad (2.7)$$

$$B(f(q, z)[P_1, \dots, P_l]) = S \left(\frac{f(q, z)}{1 - q z^{-1} P_l / P_l} [q^{-1} z P_l, P_1, P_2, \dots, P_{l-1}] \right). \quad (2.8)$$

From the definition of the space V_l , it follows that $S(q^{-1} z P_l / P_1) = z S(P_l / P_1)$ has the form $z q^n$ ($n \geq 0$) or q^m ($m < 0$). In the above, the factors $S(1 - q^{-1} z P_l / P_1)^{-1}$ and $S(1 - q z^{-1} P_l / P_l)^{-1}$ are to be understood as power series expansions in q, z .

It is immediate to check that

Lemma 2.2. *The operators $A, B : V_l \rightarrow V_l$ are well defined.*

With the aid of A, B , the recursion equations (2.4) can be recast into the following form.

Proposition 2.3. *For $N = 0, 1, \dots$, we have $\chi_{k,l;\mathbf{b}}^{(N)}(q, z) \in V_l$. Moreover*

$$\chi_{k,l;\mathbf{b}}^{(N+1)}(q, z) = (A + B) \chi_{k,l;\mathbf{b}}^{(N)}(q, z).$$

Proof. Let $\xi(q, z; \mathbf{b}, k) = f(q, z)[P_1, \dots, P_l]$. Then using (2.5), we obtain:

$$\begin{aligned} & \sum_{i=0}^{b_0} z^i \xi(q, qz; b_1 - i, \dots, b_{l-2} - i, k - i, k) \\ &= S \left(f(q, z) P_1^{b_1} P_2^{b_2 - b_1} \dots P_{l-1}^{b_{l-1} - b_{l-2}} \sum_{i=0}^{b_0} (q^{-1} z P_l / P_1)^i \right) \\ &= S \left(f(q, z) P_1^{b_1} P_2^{b_2 - b_1} \dots P_{l-1}^{b_{l-1} - b_{l-2}} \left(\frac{1}{1 - q^{-1} z P_l / P_1} + \frac{(q^{-1} z P_l / P_1)^{b_0}}{1 - q z^{-1} P_l / P_l} \right) \right) \\ &= (A + B) \xi(q, z; \mathbf{b}, k). \end{aligned}$$

The assertion follows from linearity of operators A, B and Lemma 2.1. $\square$

Note that for $N = 0$ we can write

$$\chi_{k,l;\mathbf{b}}^{(0)}(q, z) = 1 = [1, \dots, 1] \in V_l.$$

Let us denote the function $[1, \dots, 1] \in V_l$ by v_{ini} . We then obtain the following expression for the characters.

Corollary 2.4. *The characters of (k, l) -configurations are given by*

$$\chi_{k,l;\mathbf{b}}^{(N)}(q, z) = (A + B)^N v_{\text{ini}}, \quad N = 0, 1, \dots \quad (2.9)$$

Remark. The operators A and B do not commute. It would be interesting to describe explicitly the algebra generated by A, B .

2.3. Simplifications in the case $N = \infty$. The tautological formula (2.9) gives the character as a sum over all monomials in A, B of degree N . Here and below, by a *monomial* we mean an ordered composition of operators of A, B , and we assign the degree to monomials by setting $\deg A = \deg B = 1$. This expansion *a priori* comprises 2^N terms, so the number of terms grows exponentially with N . However we will see later (Section 2.4) that, at least in the limit $N \rightarrow \infty$, there are many cancellations and only polynomially many terms survive.

From the definition of the characters (2.3), it is obvious that

$$\chi_{k,l;\mathbf{b}}^{(\infty)}(q, z) = \lim_{N \rightarrow \infty} \chi_{k,l;\mathbf{b}}^{(N)}(q, z)$$

exists as a formal power series in q, z . In the limit, the recursion equations (2.4) turn into difference equations.

Proposition 2.5. *The characters $\chi_{k,l;\mathbf{b}}^{(\infty)}(q, z)$ are the unique set of formal power series in q, z satisfying the difference equation*

$$\chi_{k,l;(b_0, \dots, b_{l-2})}^{(\infty)}(q, z) = \sum_{i=0}^{b_0} z^i \chi_{k,l;(b_1-i, b_2-i, \dots, b_{l-2}-i, k-i)}^{(\infty)}(q, qz), \quad (2.10)$$

and the initial condition $\chi_{k,l;\mathbf{b}}^{(\infty)}(q, 0) = 1$.

Proof. We only need to show the uniqueness of the solution of (2.10) with a given initial condition.

The difference equation (2.10) has the form

$$y(z) = M(z)y(qz), \quad (2.11)$$

where $y(z)$ is a vector whose components are labeled by $\mathbf{b}$ satisfying (2.2), and $M(z)$ is a matrix whose non-zero entries are z^i with some i . If we expand $y(z), M(z)$ as

$$y(z) = \sum_{n \geq 0} y_n z^n, \quad M(z) = \sum_{n \geq 0} M_n z^n,$$

then (2.11) reduces to the relations

$$y_n = q^n M_0 y_n + w_n \quad (n \geq 0),$$

where w_n signifies terms which contain y_m only with $m < n$. For $n > 0$, $1 - q^n M_0$ has an inverse matrix as a formal power series in q . Hence y_n is uniquely determined by y_0 . This proves the proposition. $\square$

For the study of the difference equations (2.10), it is helpful to first solve the much simpler equation

$$Av_\infty = v_\infty, \quad v_\infty|_{z=0} = v_{\text{ini}}. \quad (2.12)$$

Notice that for each n we have

$$A^n v_{\text{ini}} = \frac{1}{(z)_n} v_{\text{ini}}, \quad (2.13)$$

where $(u)_n = \prod_{i=0}^{n-1} (1 - q^i u)$. Therefore the unique solution of (2.12) is given by

$$v_\infty = \lim_{N \rightarrow \infty} A^N v_{\text{ini}} = \frac{1}{(z)_\infty} v_{\text{ini}} = \frac{[1, \dots, 1]}{(z)_\infty} \in V_l. \quad (2.14)$$

Lemma 2.6. *We have*

$$\begin{aligned} \lim_{N \rightarrow \infty} (A + B)^N v_{\text{ini}} &= v_\infty + \sum_M MBv_\infty \\ &= (1 + B + AB + B^2 + A^2B + AB^2 + BAB + B^3 + \dots) v_\infty, \end{aligned}$$

where the sum is over all monomials M in A, B .

Proof. The lemma will follow if we show that, for any sequence $C_1, \dots, C_N$ of operators A, B , $\lim_{N \rightarrow \infty} C_1 \cdots C_N v_{\text{ini}} = 0$ holds unless $C_i = A$ for all sufficiently large i .

Let us verify this statement. In view of (2.13), it suffices to show that for any C_i we have $\lim_{N \rightarrow \infty} C_1 \cdots C_N B v_{\text{ini}} = 0$. The vector part of $B v_{\text{ini}}$ is $[z, 1, \dots, 1]$. Hence, for any monomial M'' of degree $l - 1$, we have $M'' B v_{\text{ini}} = f[P_1, \dots, P_l]$, where f is a power series and each P_i has degree 1 in z . Applying further a monomial M' of degree d , each component of the vector part of $M' M'' B v_{\text{ini}}$ acquires a power q^d due to the shift S . Therefore this expression vanishes when $d \rightarrow \infty$. $\square$

2.4. Main result. Let M, M' be monomials in A, B . We define an equivalence relation $M \sim M'$ if and only if there exist $r, r' \geq 0$ such that $MA^r = M'A^{r'}$. Lemma 2.6 is rephrased as

$$\chi_{k,l;\mathbf{b}}^{(\infty)}(q, z) = \sum_M M v_\infty, \quad (2.15)$$

where the sum ranges over all possible equivalence classes of monomials.

We say that $M = C_1 \cdots C_m$ is a *good monomial* if the following condition is satisfied.

$$\text{For all } i, C_i = A \text{ implies } C_{i+l-1} = A. \quad (2.16)$$

The notion of a good monomial makes sense also for its equivalence classes.

Example. Here is the list of all good monomials up to equivalence for $l = 2, 3, 4$.

$$\begin{aligned} l = 2 : M &= B^m \quad (m \geq 0), \\ l = 3 : M &= B^m (AB)^n \quad (m, n \geq 0), \\ l = 4 : M &= B^m (AB^2)^n (A^2 B)^p \quad (m, n, p \geq 0), \\ &B^m (BAB)^n (A^2 B)^p \quad (m \geq -1, n \geq 1, p \geq 0). \end{aligned}$$

We are now in a position to formulate our main result.

Theorem 2.7. *The character (2.15) is given by*

$$\chi_{k,l;\mathbf{b}}^{(\infty)}(q, z) = \sum_M^* M v_\infty, \quad (2.17)$$

where the sum is taken over all the equivalence classes of good monomials. Moreover each term in the sum has the form

$$\frac{(-1)^\alpha q^\beta z^\gamma}{(q)_{t_2} \cdots (q)_{t_l} (q^{t_2 + \cdots + t_l} z)_\infty},$$

with some non-negative integers $\alpha, \beta, \gamma, t_2, \dots, t_l$.

Theorem 2.7 is proved in Sections 4, 5.

Compared to formula (2.15), Theorem 2.7 provides a big simplification because good monomials are much smaller in number among all monomials. In Section 5, we give a parameterization of good monomials for general l and describe the corresponding data $\alpha, \beta, \gamma, t_2, \dots, t_l$ in combinatorial terms. Below we write down the formula (2.17) explicitly in the cases $l = 2, 3$.

Example. The case $l = 2$:

$$\begin{aligned} \chi_{k,2;b_0}^{(\infty)}(q, z) &= \sum_{n \geq 0} (-1)^{n+1} q^{n^2 k + n b_0 + 3n(n+1)/2} z^{n k + b_0 + n + 1} \frac{1}{(q)_n (q^n z)_\infty} \\ &+ \sum_{n \geq 0} (-1)^n q^{n^2 k - n b_0 + n(3n-1)/2} z^{n k + n} \frac{1}{(q)_n (q^n z)_\infty}. \end{aligned}$$

This formula has been obtained in [FL].

Example. The case $l = 3$: Set

$$g_{m,n} = q^{(3k+5)m^2/2 + (3k+5)mn + 2(k+2)n^2} z^{(k+1)(m+n)}.$$

We retain the notation $[P_1, P_2, P_3] = P_1^{b_0} P_2^{b_1 - b_0} P_3^{k - b_1}$. Then

$$\chi_{k,3;b_0,b_1}^{(\infty)}(q, z) = \sum_{m,n \geq 0} f_{m,n},$$

where $f_{m,n} = B^m(AB)^n v_\infty$ is given as follows.

$$\begin{aligned}
f_{3m,2n-1} &= (-1)^m \frac{q^{-3(k+1)m/2-(k+2)n}}{(q)_m (q)_{m+2n-1} (q^{2m+2n-1} z)_\infty} [1, q^m, q^{-m-2n+1} z^{-1}] g_{m,n}, \\
f_{3m,2n} &= (-1)^m \frac{q^{-(3k+1)m/2-kn}}{(q)_m (q)_{m+2n} (q^{2m+2n} z)_\infty} [1, q^m, q^{2m+2n}] g_{m,n}, \\
f_{3m+1,2n-1} &= (-1)^m \frac{q^{-(3k+1)m/2-(2k+1)n}}{(q)_m (q)_{m+2n} (q^{2m+2n} z)_\infty} [1, q^{m+2n}, q^{2m+2n}] g_{m,n}, \\
f_{3m+1,2n} &= (-1)^{m+1} \frac{q^{(3k+5)m/2+(2k+3)n} z^{k+1}}{(q)_m (q)_{m+2n} (q^{2m+2n} z)_\infty} [1, q^{-2m-2n} z^{-1}, q^{-m-2n} z^{-1}] g_{m,n}, \\
f_{3m+2,2n-1} &= (-1)^{m+1} \frac{q^{(3k+5)m/2+(k+2)n} z^{k+1}}{(q)_m (q)_{m+2n} (q^{2m+2n} z)_\infty} [1, q^{-2m-2n} z^{-1}, q^{-m} z^{-1}] g_{m,n}, \\
f_{3m+2,2n} &= (-1)^{m+1} \frac{q^{(3k+7)m/2+(k+4)n+1} z^{k+1}}{(q)_m (q)_{m+2n+1} (q^{2m+2n+1} z)_\infty} [1, q^{m+2n+1}, q^{-m} z^{-1}] g_{m,n}.
\end{aligned}$$

2.5. Extremal configurations. Before embarking upon the proof of Theorem 2.7, let us make a remark about the “extremal” (k, l) -configuration associated with a given monomial in (2.15).

Recall that the character (2.5) of a “segment” is written as a sum of contributions from extremal points. The same principle can be applied to write the character of a set of integer points inside any convex “polytope” in a higher dimensional space $\mathbf{R}^M$. We expect that the result is a sum of rational functions over the vertices (i.e., the extremal points) of the polytope, where each term represents the character of the cone corresponding to that vertex, see also Lemmas 3.1, 3.2 in [JMM]. We will not discuss in how much generality this assertion is valid. For a closely related discussion, see [PKh], Proposition 2.

Let us study this idea in our problem, where the set of integer points is $\mathcal{C}_{k,l;\mathbf{b}}^{(N)}$ and $M = N$. Take the simplest case $l = 2$. We consider the generic situation $0 < b_0 < k$.

For $N = 1$, the polytope is a segment $\{x_0; 0 \leq x_0 \leq b_0\}$. We have 2 vertices $a := 0$ and $b := b_0$. They correspond to Av_{ini} and Bv_{ini} , respectively.

For $N = 2$, the polytope is a quadrilateral with the 4 vertices $aa := (0, 0)$, $ab := (0, k)$, $ba := (b_0, 0)$, $bb := (b_0, k - b_0)$. They corresponds to the terms AAv_{ini} , etc.. We can classify these vertices in two groups, aa and ab with $x_0 = 0$ and ba and bb with $x_0 = b_0$. These are the vertices of the sections of $\mathcal{C}_{b_0}^{(2)}$ by the hyperplanes $x_0 = 0$ and $x_0 = b_0$. There exist natural bijections between these sections and $\mathcal{C}_k^{(1)}$ and $\mathcal{C}_{k-b_0}^{(1)}$, respectively. The vertex ab corresponds to a bad monomial AB . It turns out that in the limit $N \rightarrow \infty$ this vertex drops from the summation for the character formula. However, for $N = 2$, we cannot drop it.

For $N = 3$, the number of vertices is not equal to 8 but is 7: This is because the hyperplane section $x_0 = 0$ is isomorphic to the degenerate polytope spanned by the integer points $\mathcal{C}_0^{(2)}$, which is a triangle. Both aba and abb correspond to a single vertex $(0, k, 0)$. The corresponding monomials ABA and ABB are bad, and two terms

$ABAv_{\text{ini}}$ and $ABBv_{\text{ini}}$ cancel each other. The degree 2 bad monomial AB finds a cancellation counterpart ABB in the sense $ABv_{\infty} = ABAv_{\infty} = -ABBv_{\infty}$.

In Sections 3 and 4, we will show that such a cancellation always takes place in pairs.

In the rest of this section we describe the extremal points corresponding to good monomials. In Section 5 the character formula is written as a sum over such vertices.

To a monomial $M = C_1 \cdots C_N$, we associate a vertex $(x_0, \dots, x_{N-1}) \in \mathcal{C}_{k,l;\mathbf{b}}^{(N)}$ by the following inductive procedure. If $C_1 = A$ we set $x_0 = 0$. The section of $\mathcal{C}_{k,l;\mathbf{b}}^{(N)}$ by the hyperplane $x_0 = 0$ is isomorphic to $\mathcal{C}_{k,l;a(\mathbf{b})}^{(N-1)}$ where $a(\mathbf{b}) = (b_1, \dots, b_{l-2}, k)$. We set $(x_1, \dots, x_{N-1})$ to be the vertex corresponding to $C_2 \cdots C_N$ in $\mathcal{C}_{k,l;a(\mathbf{b})}^{(N-1)}$. If $C_1 = B$ we set $x_0 = b_0$. The section of $\mathcal{C}_{k,l;\mathbf{b}}^{(N)}$ by the hyperplane $x_0 = b_0$ is isomorphic to $\mathcal{C}_{k,l;b(\mathbf{b})}^{(N-1)}$ where $b(\mathbf{b}) = (b_1 - b_0, \dots, b_{l-2} - b_0, k - b_0)$. We set $(x_1, \dots, x_{N-1})$ to be the vertex corresponding to $C_2 \cdots C_{N-1}$ in $\mathcal{C}_{k,l;b(\mathbf{b})}^{(N-1)}$. If a vertex corresponding to M is $(x_0, \dots, x_{N-1})$, the vector part of the simple function $M(v_{\text{ini}}) = C_1 \cdots C_N(v_{\text{ini}})$ is equal to

$$P_1^{b_0} P_2^{b_1 - b_0} \cdots P_l^{k - b_{l-2}} = \prod_{i=0}^{N-1} (q^i z)^{x_i}. \quad (2.18)$$

We rephrase the above procedure as follows.

Consider an l -dimensional lattice Λ generated by formal symbols $b_0, \dots, b_{l-2}, k$. We consider $\mathbf{Z}$ linear mappings M_a, M_b on Λ given by

$$\begin{aligned} M_a(b_i) &= \begin{cases} b_{i+1} & \text{if } 0 \leq i \leq l-3; \\ k & \text{if } i = l-2, \end{cases} \\ M_a(k) &= k, \\ M_b(b_i) &= \begin{cases} b_{i+1} - b_0 & \text{if } 0 \leq i \leq l-3; \\ k - b_0 & \text{if } i = l-2, \end{cases} \\ M_b(k) &= k. \end{aligned}$$

For a given monomial $M = C_1 \cdots C_N$ we set

$$c_i = \begin{cases} a & \text{if } C_i = A; \\ b & \text{if } C_i = B. \end{cases}$$

Then we have

$$x_i = \begin{cases} 0 & \text{if } C_{i+1} = A; \\ M_{c_1} \circ \cdots \circ M_{c_i}(b_0) & \text{if } C_{i+1} = B. \end{cases} \quad (2.19)$$

If M is a good monomial, the above procedure is simplified. First we prepare some notation and a few lemmas.

Define $\iota_{i,j} \in \Lambda$ for $i, j \in \mathbf{Z}$ by

$$\iota_{i,j} = b_{\bar{i}-2} - b_{\bar{j}-2} + k\delta(\bar{i} \leq \bar{j}) \in \Lambda, \quad (2.20)$$

where $b_{-1} = 0$, $\bar{i}$ and $\bar{j}$ are such that $1 \leq \bar{i}, \bar{j} \leq l$, $i \equiv \bar{i} \pmod{l}$, $j \equiv \bar{j} \pmod{l}$, and

$$\delta(*) = \begin{cases} 1 & \text{if } * \text{ is true;} \\ 0 & \text{otherwise.} \end{cases}$$

Note that $\iota_{2,1} = b_0 \in \Lambda$ and $\iota_{a,b} \neq 0$ in Λ . In the computation of x_i in (2.19), we start from $\iota_{2,1}$, and apply c_i, c_{i-1} , etc., successively. We will prove that if M is good we will never get 0 in this process.

Let us start with

Lemma 2.8. *The actions of M_a and M_b on Λ satisfy*

$$\begin{aligned} M_a(\iota_{i,j}) &= \begin{cases} \iota_{i+1-\delta_{i,1}, j+1-\delta_{j,1}} & \text{if } (\bar{i}, \bar{j}) \neq (1, l); \\ 0 & \text{if } (\bar{i}, \bar{j}) = (1, l), \end{cases} \\ M_b(\iota_{i,j}) &= \iota_{i+1, j+1}. \end{aligned}$$

Proof. The proof is only case checking. □

Lemma 2.9. *Let $M = C_1 \cdots C_N$ be a good monomial, and $C_i = B$. We have $M_{c_1} \circ \cdots \circ M_{c_{i-1}}(\iota_{2,1}) \neq 0$.*

Proof. Consider the mapping $a, b : \{1, \dots, l\} \rightarrow \{1, \dots, l\}$ given by

$$\begin{aligned} a(j) &= \begin{cases} j+1 & \text{if } j \neq 1, l; \\ 1 & \text{otherwise,} \end{cases} \\ b(j) &= \begin{cases} j+1 & \text{if } j \neq l; \\ 1 & \text{if } j = l. \end{cases} \end{aligned}$$

We have $M_a(\iota_{j,j'}) = \iota_{a(j), a(j')}$ unless $(j, j') = (1, l)$, and $M_b(\iota_{j,j'}) = \iota_{b(j), b(j')}$. Suppose that $C_i = B$. The trajectory of 1 by the successive mappings c_x , where x changes its values as $i-1, i-2$, etc., stays at 1 exactly at the following values of x . The trajectory starts at 1 for $x = i$. It stays at 1 while $C_x = A$. It changes to 2 when $C_x = B$ for the first time after $x = i$. Suppose that this is when $x = i_1$. Then, the trajectory comes back to 1 when $x = i_1 - l + 1$. Since M is good, we have $C_{i_1-l+1} = B$. It stays at 1 while $C_x = A$, and changes to 2 when $C_x = B$ for the first time after $x = i_1 - l + 1$. This pattern repeats. In particular, the trajectory comes to l only if the next C_x is equal to B .

On the trajectory of $\iota_{2,1}$ at $x = i$, 0 appears only if M_a applies to $\iota_{1,l}$. However, since the trajectory of 1 at $x = i$ comes to l only if the next mapping is M_b , this never occurs. □

Fix a good monomial $M = C_1 \cdots C_N$. We define C_i , $i \leq 0$, by setting $C_i = B$. For $i \geq 1$ such that $C_i = B$ we define

$$\begin{aligned} \gamma_i &= c_1 \circ \cdots \circ c_{i-1}(2), \\ \gamma'_i &= c_1 \circ \cdots \circ c_{i-1}(1). \end{aligned}$$

Because of Lemma 2.8 and 2.9, we can rewrite (2.19) into $x_i = \iota_{\gamma_{i+1}, \gamma'_{i+1}}$ for i such that $C_{i+1} = B$. We define i_- to be the largest integer satisfying $i_- < i$ and $C_{i_-} = B$. Then we have the recursion relation for γ_i and γ'_i :

$$\gamma'_i = \gamma_{i_-}, \quad (2.21)$$

$$\gamma_i = \begin{cases} i+1 & \text{if } i \leq l-1; \\ \gamma'_{i-l+1} & \text{if } i \geq l. \end{cases} \quad (2.22)$$

The second relation makes sense because $C_{i-l+1} = B$, which follows from $C_i = B$ and the assumption that M is good.

Therefore, we have

Corollary 2.10. *If M is a good monomial and $C_{i+1} = B$ then*

$$x_i = \iota_{\gamma_{i+1}, \gamma'_{i+1}},$$

where $\gamma_{i+1}, \gamma'_{i+1}$ are given by (2.21), (2.22).

We defined the equivalence of monomials by $M \sim MA$. If $(x_0, \dots, x_{N-1})$ is the extremal configuration for M then the extremal configuration for MA is $(x_0, \dots, x_{N-1}, 0)$. We define the equivalence of configurations so that $(x_0, \dots, x_{N-1}) \sim (x_0, \dots, x_{N-1}, 0)$. The extremal configuration corresponding to an equivalence class of monomials is well-defined.

Proposition 2.11. *Suppose $M = C_1 \cdots C_N$ is a good monomial. If a monomial M' gives a configuration equivalent to that of M , then M' is equivalent to M .*

Proof. From Lemma 2.9, if M' is good its equivalence class is uniquely determined from the corresponding configuration. Suppose $M' = C'_1 \cdots C'_{N'}$ is bad. There exists $i \geq 1$ such that $C_i = A$, $C'_i = B$, and $C_j = C'_j$ for $j < i$. Since $C_i = A$ and the configurations corresponding to M, M' are equivalent, we have $c_1 \circ \cdots \circ c_{i-1}(\iota_{21}) = 0$. However, this is impossible because $C_1 \cdots C_{i-1}$ is good, and by the same argument as in Lemma 2.9, we can show that if $c_j \circ \cdots \circ c_{i-1}(1) = l$ then C_{j-1} is necessarily B . $\square$

In Section 5, we will show that if M is bad, there exists another bad monomial M' such that $(M + M')v_\infty = 0$. Such a pair of (equivalence classes of) monomials will be called a cancellation pair. In particular, we will show (Proposition 3.5) that the extremal configurations for a cancellation pair are identical.

3. CANCELLATION PAIRS

In this section we give a combinatorial preparation which will be used to prove Theorem 2.7.

3.1. Summation graph. Let M be a monomial in A, B of degree m . It acts on a vector $f[P_1, \dots, P_l]$ as

$$M(f[P_1, \dots, P_l]) = S^m(f)\varphi_M(P_1, \dots, P_l)[\tilde{P}_1, \dots, \tilde{P}_l].$$

Here S denotes the q -shift (2.6), and $\varphi_M(P_1, \dots, P_l)$ is some scalar function determined from $[P_1, \dots, P_l]$. The vector part $[\tilde{P}_1, \dots, \tilde{P}_l]$ has the structure

$$[q^{r_1} z^{s_1} S^m(P_{\sigma_M(1)}), \dots, q^{r_l} z^{s_l} S^m(P_{\sigma_M(l)})] \quad (r_i, s_i \in \mathbb{Z}),$$

where $\sigma_M : \{1, \dots, l\} \rightarrow \{1, \dots, l\}$ is some map which is not necessarily bijective. We have

$$\sigma_A(1) = 1, \quad \sigma_A(i) = i - 1 \quad (2 \leq i \leq l), \quad (3.1)$$

$$\sigma_B(1) = l, \quad \sigma_B(i) = i - 1 \quad (2 \leq i \leq l). \quad (3.2)$$

Moreover,

$$\sigma_{MM'} = \sigma_{M'} \circ \sigma_M. \quad (3.3)$$

Note that the map σ_M is uniquely specified by (3.1)–(3.3).

Let us classify monomials according to the subset

$$\{\sigma_M(1), \dots, \sigma_M(l)\} \subset \{1, \dots, l\}.$$

For that purpose it is convenient to introduce an oriented graph G , which we call the *summation graph*. By definition, the vertices of G are nonempty subsets of $\{1, 2, \dots, l\}$. We represent a vertex I by an array of length l , whose i -th component is $\bullet$ or $\circ$ depending on whether $i \in I$ or not. For example, $[\bullet \circ \circ \bullet \bullet]$ means $I = \{1, 4, 5\}$ with $l = 5$. Each vertex is a source of two arrows, arrow A and arrow B . These arrows connect vertices by the following rule.

$$[\bullet * J] \xrightarrow{A} [\bullet J \circ], \quad (3.4)$$

$$[\circ J'] \xrightarrow{A} [J' \circ]. \quad (3.5)$$

$$[* J'] \xrightarrow{B} [J' *]. \quad (3.6)$$

Here $*$ = $\bullet, \circ$, and J, J' are sequences of $\bullet, \circ$ of length $l - 2$ or $l - 1$, respectively. There are only two arrows whose source and sink coincide,

$$\begin{aligned} [\overbrace{\bullet \dots \bullet}^l] &\xrightarrow{B} [\overbrace{\bullet \dots \bullet}^l], \\ [\bullet \overbrace{\circ \dots \circ}^{l-1}] &\xrightarrow{A} [\bullet \overbrace{\circ \dots \circ}^{l-1}]. \end{aligned}$$

The summation graph G in the case $l = 3$ is shown in fig.1.

We choose the vertex $I_{top} = [\bullet \dots \bullet]$ with all $\bullet$, and fix it as a reference vertex. Given a monomial $M = C_1 C_2 \dots C_m$ ($C_i \in \{A, B\}$), let us follow the arrows $C_1, C_2 \dots$ starting from I_{top} . We call the resulting sequence of vertices $\{I^{(i)}\}$,


$$I_{top} = I^{(0)} \xrightarrow{C_1} I^{(1)} \xrightarrow{C_2} \dots \xrightarrow{C_m} I^{(m)}, \quad (3.7)$$

the *path associated with M* . We say that the path (or, by abuse of language, the monomial M) *passes through* the vertices $I^{(i)}$. With the rule above, it is easy to see the following.

Lemma 3.1. *The subset $\{\sigma_M(1), \dots, \sigma_M(l)\}$ is given by the last vertex $I^{(m)}$ in the path.*

For example, if $l = 5$ and $M = A^2 B^2 A$, then

$$\begin{aligned} [\bullet \bullet \bullet \bullet \bullet] &\xrightarrow{A} [\bullet \bullet \bullet \bullet \circ] \xrightarrow{A} [\bullet \bullet \bullet \circ \circ] \\ &\xrightarrow{B} [\bullet \bullet \circ \circ \bullet] \xrightarrow{B} [\bullet \circ \circ \bullet \bullet] \xrightarrow{A} [\bullet \circ \bullet \bullet \circ], \end{aligned} \quad (3.8)$$


 FIGURE 1. Summation graph ($l = 3$).

hence $\{\sigma_M(1), \dots, \sigma_M(5)\} = \{1, 3, 4\}$. Notice that in this procedure the monomial should be read from *left to right* because of (3.3).

We say that a vertex I is *good* if $1 \in I$, or equivalently, if the first entry of I is $\bullet$. The notion of good monomials (2.16) has the following interpretation.

Lemma 3.2. *A monomial is good if and only if it passes through only good vertices.*

Proof. We retain the notation of (3.7). If $C_i = A$, then there is a $\circ$ in the l -th component of $I^{(i)}$. This $\circ$ will then move to the left by one step with each A or B arrow, until it reaches the second component in the array. Hence the first component of $I^{(i+l-1)}$ is $\bullet$ if and only if $C_{i+l-1} = A$. $\square$

3.2. Marked vertices. In dealing with paths associated with monomials, it is often convenient to keep track of the move of the last entry of I_{top} . Let us introduce paths with marked vertices. A marked vertex is a pair (I, j) of vertex I and an element $j \in I$. The element j is represented by the symbol $\times$ in place of $\bullet$. We extend the definition of the A, B arrows to marked vertices as follows.

$$[\circ J] \xrightarrow{A} [J \circ], \quad (3.9)$$

$$[\bullet \bullet J] \xrightarrow{A} [\bullet J \circ], \quad (3.10)$$

$$[\bullet \circ J] \xrightarrow{A} [\bullet J \circ], \quad (3.11)$$

$$[\bullet \times J] \xrightarrow{A} [\bullet J \circ], \quad (3.12)$$

$$[\times \bullet J] \xrightarrow{A} [\bullet J \circ], \quad (3.13)$$

$$[\times \circ J] \xrightarrow{A} [\times J \circ], \quad (3.14)$$

$$[\bullet J] \xrightarrow{B} [J \bullet], \quad (3.15)$$

$$[\circ J] \xrightarrow{B} [J \circ], \quad (3.16)$$

$$[\times J] \xrightarrow{B} [J \times]. \quad (3.17)$$

Here J in (3.9–3.11) and (3.15–3.16) contain a $\times$, and others not. Note that in (3.12–3.13) $\times$ is lost in the RHS.

Starting with $[\bullet \cdots \bullet \times]$ we obtain a path P passing through marked or unmarked vertices similar to (3.7). We call it the *marked path* associated with M . For example, if $l = 5$ and $M = AB^2A$, then

$$\begin{aligned} [\bullet \bullet \bullet \bullet \times] &\xrightarrow{A} [\bullet \bullet \bullet \times \circ] \xrightarrow{B} [\bullet \bullet \times \circ \bullet] \\ &\xrightarrow{B} [\bullet \times \circ \bullet \bullet] \xrightarrow{A} [\bullet \circ \bullet \bullet \circ]. \end{aligned} \quad (3.18)$$

The rules (3.9–3.17) are so designed that, replacing $\times$ with either $\bullet$ or $\circ$, we get back to the original rule (3.4)–(3.6) for unmarked vertices. The marked path P provides a one-to-one correspondence between the unmarked path associated with a monomial M and that of AM . Indeed, the former is recovered by replacing $\times$ with $\bullet$ in P . The latter is obtained by first supplementing P with a leftmost arrow $[\bullet \cdots \bullet] \xrightarrow{A}$ and then replacing $\times$ with $\circ$. For instance, the second procedure applied to the example (3.18) gives (3.8) for $AM = A^2B^2A$. This correspondence will be used extensively in the sequel.

3.3. Cancellation pair. Henceforth we regard an equivalence class of monomials as a monomial $M = C_1 \cdots C_m$ such that m is sufficiently large and $C_i = A$ for sufficiently large i .

Consider the marked path P associated with a good monomial M . If P does not pass through a vertex which has $\times$ as the first entry, then AM is good. Otherwise, P contains one of the following arrows.

- (i) $[\times \bullet J] \xrightarrow{A} [\bullet J \circ],$
- (ii) $[\times \bullet J] \xrightarrow{B} [\bullet J \times],$
- (iii) $[\times \circ J] \xrightarrow{A} [\times J \circ],$
- (iv) $[\times \circ J] \xrightarrow{B} [\circ J \times].$

The case (iv) does not occur since M is good. If (iii) takes place for some arrow $C_j = A$, the J must contain $\bullet$. This is because A does not bring $\times$ to the first column, and when B does so, it brings $\bullet$ back to the last column. Thus, we can find a $j' > j$ such that $C_{j'}$ is either (i) or (ii). Among the arrows of type (i) or (ii), let C_i be such that the index i is the largest. We call C_i a *cancellation A-arrow* in the case (i), and a *cancellation B-arrow* in the case (ii).

Denote by $\mathcal{G}$ the set of equivalence classes of good monomials. We have shown that it is a disjoint union of three subsets,

$$\mathcal{G} = \mathcal{G}_G \cup \mathcal{G}_A \cup \mathcal{G}_B.$$

Here $\mathcal{G}_G = \{M \in \mathcal{G} \mid AM \in \mathcal{G}\}$, and $\mathcal{G}_A$ (resp. $\mathcal{G}_B$) is the set of good monomials which has a cancellation A-arrow (resp. B-arrow).

Lemma 3.3. *There is a bijection between $\mathcal{G}_A$ and $\mathcal{G}_B$.*

Proof. Let $M = C_1 \cdots C_m = NAK \in \mathcal{G}_A$ where $N = C_1 \cdots C_{i-1}$ and $K = C_{i+1} \cdots C_m$. Suppose that $C_i = A$ is the cancellation A-arrow. Then, K is a good monomial, and

the path after C_i has the form

$$[\times \bullet J] \xrightarrow{A} [\bullet J \circ] \xrightarrow{C_{i+1}} \dots \xrightarrow{C_{i+l-2}} [\bullet \circ J'] \xrightarrow{A} [\bullet J' \circ] \xrightarrow{C_{i+l}} \dots$$

Replacing the cancellation A -arrow by B , the path associated with NBK after B becomes

$$[\times \bullet J] \xrightarrow{B} [\bullet J \times] \xrightarrow{C_{i+1}} \dots \xrightarrow{C_{i+l-2}} [\bullet \times J'] \xrightarrow{A} [\bullet J' \circ] \xrightarrow{C_{i+l}} \dots$$

This shows that the B in NBK is the cancellation B -arrow.

Hence we obtain a map from $\mathcal{G}_A$ into $\mathcal{G}_B$. The inverse map can be constructed similarly. $\square$

We call the pair (NAK, NBK) described in the proof a *cancellation pair*.

3.4. Reduction of the Theorem. Let us turn to the proof of Theorem 2.7. Denote by $\chi^*(q, z)$ the right hand side of (2.17). We must show that $\chi^*(q, z) = \chi_{k,l;b}(q, z)$. Clearly $\chi^*(q, 0) = 1$. Hence, in view of Lemma 2.5, it is enough to verify the difference equation

$$\chi^*(q, z) = (A + B)\chi^*(q, z).$$

The left hand side can be written as

$$\sum_{M \in A\mathcal{G} \cap \mathcal{G}} Mv_\infty + \sum_{M \in B\mathcal{G} \cap \mathcal{G}} Mv_\infty,$$

because by the definition if AM' or BM' is a good monomial then M' is also a good monomial. Note further that $A\mathcal{G} \cap \mathcal{G} = A\mathcal{G}_G$ and $B\mathcal{G} \cap \mathcal{G} = B\mathcal{G}$.

The right hand side gives

$$\sum_{M' \in \mathcal{G}_G} AM'v_\infty + \sum_{M' \in \mathcal{G}_A} AM'v_\infty + \sum_{M' \in \mathcal{G}_B} AM'v_\infty + \sum_{M' \in \mathcal{G}} BM'v_\infty.$$

Combining these relations with Lemma 3.3, we see that the proof of the first statement of Theorem 2.7 reduces to the following statement.

Theorem 3.4. *For any cancellation pair $(M, M') = (NAK, NBK)$, we have*

$$AN(A + B)Kv_\infty = 0. \quad (3.19)$$

The proof of Theorem 3.4 will be given in the next section.

Here we give a proof of the following statement.

Proposition 3.5. *Suppose that (NAK, NBK) is a cancellation pair. The extremal configurations for the pair of monomials $(ANAK, ANBK)$ are identical.*

Proof. Suppose that $AN = C_1 \dots C_{i-1}$, and $K = C_{i+1} \dots C_M$. We are to show that x_j and x'_j corresponding to $ANAK$ and $ANBK$, respectively, are in fact equal. There are four cases (i) $j \leq i - 1$, (ii) $j = i - 1$, (iii) $j \geq i$ and $C_{j+1} = A$, and (iv) $j \geq i$ and $C_{j+1} = B$.

The statement is clear if $j \leq i - 2$ because the part AN is common to $ANAK$ and $ANBK$.

For $j = i - 1$ since $x_{i-1} = 0$, we must show that $x'_{i-1} = M_{c_1} \circ \dots \circ M_{c_{i-1}}(\iota_{2,1}) = 0$. Consider the marked path $\{I^{(a)}\}$ corresponding to the monomial $ANBK$. We have

$I^{(0)} = [\bullet \cdots \bullet \times]$. Let $1 \leq \eta_a, \eta'_a \leq l$ be such that the position of $\times$ in the marked vertex $I^{(a)}$ is η_a , and that of the first $\bullet$ we find when we move from $\times$ to the right cyclically is η'_a . It is easy to see that $\eta_a = c_{a+2} \circ \cdots \circ c_{i-1}(1)$ and $\eta'_a = c_{a+2} \circ \cdots \circ c_{i-1}(2)$. We have $(\eta'_{i-2}, \eta_{i-2}) = (2, 1)$ and $(\eta'_0, \eta_0) = (1, l)$. Therefore, $M_{c_1} \circ \cdots \circ M_{c_{i-1}}(\iota_{2,1}) = 0$.

Now consider $j \geq i$. If $C_{j+1} = A$ then $x_j = x'_j = 0$. Take $j \geq i$ such that $C_{j+1} = B$. We will show that $M_{c_1} \circ \cdots \circ M_a \cdots \circ M_{c_j}(\iota_{2,1}) = M_{c_1} \circ \cdots \circ M_b \cdots \circ M_{c_j}(\iota_{2,1})$, where M_a in the LHS and M_b in the RHS are at the place i . Since NAK and NBK are good, $M_a(\iota_{1,l}) = 0$ never occurs on the trajectories of $\iota_{2,1}$ in these formulas. Therefore, it is enough to show $c_1 \circ \cdots \circ a \cdots \circ c_j(s) = c_1 \circ \cdots \circ b \cdots \circ c_j(s)$ for both $s = 1, 2$. Note that $a(t) = b(t)$ unless $t = 1$. On the other hand, $c_2 \circ \cdots \circ a(1) = l, c_2 \circ \cdots \circ b(1) = 1$ as shown above. Therefore, $c_1 \circ \cdots \circ a(1) = c_1 \circ \cdots \circ b(1) = 1$. This proves the statement. $\square$

4. PROOF OF THE CANCELLATION

4.1. The pairwise cancellation. Let us describe the mechanism of the cancellation (3.19). First we note a simple fact which is a corollary of the definition of A, B :

Lemma 4.1. *For any simple vector v , let*

$$Av = f[P_1, \cdots, P_l], \quad Bv = f'[P'_1, \cdots, P'_l].$$

Then

$$\begin{aligned} P_1 &= P_2, \quad P_j = P'_j \quad (2 \leq j \leq l), \\ f/f' &= -P'_2/P'_1. \end{aligned}$$

In view of this Lemma, we set

$$BNAKv_\infty = g[Q_1, \cdots, Q_l], \tag{4.1}$$

$$BNBKv_\infty = g'[Q'_1, \cdots, Q'_l], \tag{4.2}$$

$$AKv_\infty = h[R_1, \cdots, R_l], \tag{4.3}$$

$$BKv_\infty = h'[R'_1, \cdots, R'_l]. \tag{4.4}$$

From Lemma 4.1 we have

$$\begin{aligned} ANAKv_\infty &= -\frac{Q_2}{Q_1} g[Q_2, Q_2, \cdots, Q_l], \\ ANBKv_\infty &= -\frac{Q'_2}{Q'_1} g'[Q'_2, Q'_2, \cdots, Q'_l]. \end{aligned}$$

The proof of the following proposition is given in Section 4.4.

Proposition 4.2. *Set $m = \deg BN$. Then*

- (i) $Q_j = Q'_j$ for $2 \leq j \leq l$,
- (ii) $Q_2/Q_1 = q^\alpha S^m(R_2/R_1)$ and $Q'_2/Q'_1 = q^\alpha S^m(R'_2/R'_1)$, where α is an integer,
- (iii) $g/g' = S^m(h/h')$.

Assuming this proposition we give

Proof. (of Theorem 3.4) We have

$$\left(-\frac{Q_2}{Q_1}g\right) / \left(-\frac{Q'_2}{Q'_1}g'\right) = S^m \left(\frac{R_2}{R_1} \frac{R'_1}{R'_2} \frac{h}{h'}\right) = -1,$$

where in the second equality we used Lemma 4.1. Therefore

$$ANAKv_\infty + ANBKv_\infty = 0.$$

□

The rest of this section is devoted to the proof of Proposition 4.2.

4.2. Cancellation blocks. In this subsection we study the structure of the monomials BN and K when (NAK, NBK) constitutes a cancellation pair.

Let s be an integer satisfying $0 \leq s \leq l-2$. We say that a monomial of degree $l+s$

$$\mathbb{B}_s = C_1 C_2 \cdots C_{l+s}$$

is a *cancellation block* if the following two conditions are satisfied.

- (i) $C_1 = C_{s+2} = C_l = B$,
- (ii) $C_2 = \cdots = C_{s+1} = A$, $C_{l+1} = \cdots = C_{l+s} = A$.

Lemma 4.3. *Let $(M, M') = (NAK, NBK)$ be a cancellation pair. Then BN can be decomposed uniquely as a product of cancellation blocks,*

$$BN = \mathbb{B}_{s_1} \cdots \mathbb{B}_{s_p},$$

with some $p \geq 1$ and $0 \leq s_1, \dots, s_p \leq l-2$.

Proof. First note the following. Suppose a marked vertex has $\times$ at the last column. If we apply only A arrows on this vertex, we will never reach a cancellation A -arrow.

Define $s_1 \geq 0$ to be the integer such that N has the form $A^{s_1}BN_1$ with some monomial N_1 . The existence of such B in N is assured by the above remark.

We write $BN = C_1 C_2 \cdots$ with $C_i = A$ or B , so that $C_1 = B$, $C_2 = \cdots = C_{s_1+1} = A$ and $C_{s_1+2} = B$. Starting with the marked vertex $[\times \bullet \cdots \bullet]$ let us keep track of its changes under successive applications of $C_1, C_2, \dots$.

For the first $s_1 + 1$ letters BA^{s_1} we obtain

$$[\times \bullet \cdots \bullet] \xrightarrow{B} [\bullet \cdots \bullet \times] \xrightarrow{A} \cdots \xrightarrow{A} [\bullet \cdots \bullet \times \overbrace{\circ \cdots \circ}^{s_1}].$$

From the definition (2.16) of a good monomial follows that $s_1 \leq l-2$ since otherwise B cannot appear in N . Continuing with the move after BA^{s_1} and noting $C_{s_1+2} = B$, we have

$$\begin{aligned} & [\bullet \cdots \bullet \times \overbrace{\circ \cdots \circ}^{s_1}] \xrightarrow{B} [\bullet \cdots \bullet \times \overbrace{\circ \cdots \circ}^{s_1} \bullet] \\ & \xrightarrow{C_{s_1+3}} [\bullet \cdots \bullet \times \overbrace{\circ \cdots \circ}^{s_1} \bullet \bar{C}_{s_1+3}] \xrightarrow{C_{s_1+4}} \cdots \xrightarrow{C_{l-1}} [\bullet \times \overbrace{\circ \cdots \circ}^{s_1} \bullet \bar{C}_{s_1+3} \cdots \bar{C}_{l-1}], \end{aligned}$$

where $\bar{C}_i$ signifies $\circ$ if $C_i = A$, and $\bullet$ otherwise. The next letter C_l must be B , since otherwise the $\times$ will disappear by A and it would mean that $ANAK$ is a good monomial.

After that, only A (s_1 times) is allowed, and we arrive at

$$[\bullet \times \overbrace{\circ \cdots \circ}^{s_1} \bullet \bar{C}_{s_1+3} \cdots \bar{C}_{l-1}] \xrightarrow{BA^{s_1}} [\times \bullet \bar{C}_{s_1+3} \cdots \bar{C}_{l-1} \bullet \overbrace{\circ \cdots \circ}^{s_1}].$$

If the next letter is A , this is a cancellation arrow. In this case we have shown that $BN = \mathbb{B}_{s_1}$ with $\mathbb{B}_{s_1} = BA^{s_1}BC_{s_1+3} \cdots C_{l-1}BA^{s_1}$. Otherwise, $BN = \mathbb{B}_{s_1}N'$ with $N' = B \cdots$. We can now repeat the same process for N' , starting with the vertex

$$[\times \bullet \bar{C}_{s_1+3} \cdots \bar{C}_{l-1} \bullet \overbrace{\circ \cdots \circ}^{s_1}].$$

After a finite number of steps we exhaust BN and arrive at the cancellation A -arrow. $\square$

Lemma 4.4. *In the notation of Lemma 4.3, let $\mathbb{B}_{s_i} = C_1^{(i)} \cdots C_{l+s_i}^{(i)}$. Then, for each $1 \leq i \leq p-1$, one of the following conditions holds.*

- (i) $s_{i+1} = l-2$,
- (ii) $l \geq s_i + s_{i+1} + 3$ and $C_{l-s_i}^{(i+1)} = C_{l-s_i+1}^{(i+1)} = \cdots = C_{l-1}^{(i+1)} = A$.

Proof. To simplify the notation let us write $s = s_i$, $t = s_{i+1}$, $\mathbb{B}_s = \mathbb{B}_{s_i} = C_1 \cdots C_{l+s}$ and $\mathbb{B}_t = \mathbb{B}_{s_{i+1}} = D_1 \cdots D_{l+t}$. We have seen that $\mathbb{B}_s$ causes a change of vertices

$$[\times \bullet \cdots] \longrightarrow [\times \bullet \bar{C}_{s+3} \cdots \bar{C}_{l-1} \bullet \overbrace{\circ \cdots \circ}^s].$$

We now continue with $\mathbb{B}_t$. The leftmost B in $\mathbb{B}_t$ changes the right hand side to

$$[\bullet \bar{C}_{s+3} \cdots \bar{C}_{l-1} \bullet \overbrace{\circ \cdots \circ}^s \times].$$

We distinguish the three cases.

- (i) $t = l-2$,
- (ii) $l \geq s+t+3$,
- (iii) $1 \leq l-t-2 \leq s$.

The case (iii) is impossible. Indeed, if (iii) takes place, then $D_2 \cdots D_{t+1} = A^t$ changes the vertex to

$$[\bullet \overbrace{\circ \cdots \circ}^{l-t-2} \times \overbrace{\circ \cdots \circ}^t],$$

and the next letter $D_{t+2} = B$ leads to a bad vertex.

In the case (ii), A^t followed by the next $l-s-t-2$ letters $D_{t+2} \cdots D_{l-s-1}$ change the vertex to

$$[\bullet \overbrace{\circ \cdots \circ}^s \times \overbrace{\circ \cdots \circ}^t \bar{D}_{t+2} \cdots \bar{D}_{l-s-1}].$$

By the same reason as above, the next s letters $D_{l-s}, \dots, D_{l-1}$ must all be A , the case $s=0$ inclusive. The proof is over. $\square$

Lemma 4.5. *We retain the notation of Lemma 4.3. Then K has the form $\mathbb{D}_1 \cdots \mathbb{D}_r$, where $\mathbb{D}_i = \mathbb{D}'_i A^{s_p+1}$ and $\mathbb{D}'_i$ has degree $l-s_p-2$.*

Proof. Recall the definition (2.16). The proof is straightforward because NAK is a good monomial, and N contains A^{s_p} at the right end. $\square$

4.3. Change of vectors. Our next task is to study how the vectors change under the action of a cancellation block.

Lemma 4.6. *Let $M = C_1 \cdots C_l$ be an arbitrary monomial of degree l , and set*

$$M(f[P_1, \dots, P_l]) = \tilde{f}[\tilde{P}_1, \dots, \tilde{P}_l].$$

- (i) *If $C_i = A$, then $\tilde{P}_i = \tilde{P}_{i+1}$,*
- (ii) *If $C_i = B$, then $\tilde{P}_i = q^{i-1}zS^l(P_i)$.*

Proof. For each $i = 1, \dots, l$ let $C_i \cdots C_l[P_1, \dots, P_l] = f^{(i)}[P_1^{(i)}, \dots, P_l^{(i)}]$. If $C_i = A$, then $P_1^{(i)} = P_2^{(i)}$. If $C_i = B$, then $P_1^{(i)} = zS(P_l^{(i+1)}) = zS^{l-i+1}(P_i)$. Noting that $\tilde{P}_i = P_i^{(1)} = S^{j-1}(P_{i-j+1}^{(j)})$ ($i \geq j$), we obtain the assertion. $\square$

Now let $\mathbb{B}_s = C_1 \cdots C_{l+s}$ be a cancellation block of degree $l + s$. Given a vector $v = f[P_1, \dots, P_l]$, we set

$$\mathbb{B}_s f([P_1, \dots, P_l]) = \tilde{f}[\tilde{P}_1, \dots, \tilde{P}_l]. \quad (4.5)$$

Lemma 4.7. *In the notation of (4.5), the following hold for $1 \leq i \leq l$.*

- (i) *If $C_i = A$, then $\tilde{P}_i = \tilde{P}_{i+1}$,*
- (ii) *If $C_i = B$, then*

$$\tilde{P}_i = \begin{cases} q^{i-1}zS^{l+s}(P_1) & (1 \leq i \leq s+1), \\ q^{i-1}zS^{l+s}(P_{i-s}) & (s+2 \leq i \leq l). \end{cases}$$

- (iii) $\tilde{P}_i/\tilde{P}_1 = q^{s+1}S^{l+s}(P_2/P_1)$ ($2 \leq i \leq s+2$).

Proof. The assertions (i),(ii) follow from Lemma 4.6 applied to

$$A^s(f[P_1, \dots, P_l]) = f'[\overbrace{S^s(P_1), \dots, S^s(P_1)}^{s+1}, S^s(P_2), \dots, S^s(P_{l-s})].$$

Since $C_1 = C_{s+2} = B$, we have $\tilde{P}_1 = zS^{l+s}(P_1)$ and $\tilde{P}_{s+2} = q^{s+1}zS^{l+s}(P_2)$. On the other hand, $C_2 = \dots = C_{s+1} = A$ implies $\tilde{P}_2 = \dots = \tilde{P}_{s+2}$, whence follows the assertion (iii). $\square$

Let us study the scalar part $\tilde{f}$ in (4.5). Define $v^{(j)}$ inductively by $v^{(l+s+1)} = v$ and

$$v^{(j)} = C_j v^{(j+1)} = f^{(j)}[P_1^{(j)}, \dots, P_l^{(j)}].$$

We have $\tilde{f} = f^{(1)}$. In general, $f^{(j)}$ has the form

$$f^{(j)} = \frac{1}{1-g_j} S\left(\frac{1}{1-g_{j+1}}\right) \cdots S^{l+s-j}\left(\frac{1}{1-g_{l+s}}\right).$$

The factors g_j are determined as follows.

Lemma 4.8. *In the above setting, we have*

$$g_1 = q^{s+1} S^{l+s}(P_2/P_1), \quad g_j = q^{j-s-2} S^{l+s+1-j}(P_1/P_2) \quad (2 \leq j \leq s+1), \quad (4.6)$$

$$g_l = z^{-1} S^{s+1}(P_1/P_{l-s}), \quad g_j = z S^{l+s+1-j}(P_{j-s}/P_1) \quad (l+1 \leq j \leq l+s), \quad (4.7)$$

$$g_{s+2}, \dots, g_{l-1} \text{ do not depend on } P_1. \quad (4.8)$$

Proof. Let us calculate g_j 's by applying $C_{l+s}, C_{l+s-1}, \dots$ step by step. The formula (4.7) is easily obtained. After $s+1$ steps, $[P_1^{(l-1)}, \dots, P_l^{(l-1)}]$ is given by

$$[z S^{s+1}(P_{l-s}), \overbrace{S^{s+1}(P_1), \dots, S^{s+1}(P_1)}^{s+1}, S^{s+1}(P_2), \dots, S^{s+1}(P_{l-s-1})].$$

Hence during the next $l-s-2$ steps the computation of g_j involves only $P_2, \dots, P_{l-s}$. Since $C_{s+2} = B$, $[P_1^{(s+2)}, \dots, P_l^{(s+2)}]$ takes the form

$$[z S^{l-1}(P_2), *, \dots, *, \overbrace{S^{l-1}(P_1), \dots, S^{l-1}(P_1)}^{s+1}].$$

The g_j for $1 \leq j \leq s+1$ can be calculated from this. $\square$

We say that a pair of simple vectors (v, v') satisfies the *condition* $(\mathbf{C})_s$ ($0 \leq s \leq l-2$) if

$$(\mathbf{C1})_s: P_j = P'_j \quad (2 \leq j \leq l),$$

$$(\mathbf{C2})_s: P_1/P_2 = z P'_1/P'_1,$$

$$(\mathbf{C3})_s: P_{l-s} = P_{l-s+1} = \dots = P_l,$$

where $v = f[P_1, \dots, P_l]$, $v' = f'[P'_1, \dots, P'_l]$. Because of $(\mathbf{C1})_s$, these conditions are symmetric in $\{P_j\}_{j=1}^l$ and $\{P'_j\}_{j=1}^l$.

Let (v, v') be a pair satisfying $(\mathbf{C})_s$. Let $\mathbb{B}_s$ be a cancellation block, and set

$$v = f[P_1, \dots, P_l],$$

$$v' = f'[P'_1, \dots, P'_l],$$

$$\mathbb{B}_s v = \tilde{f}[\tilde{P}_1, \dots, \tilde{P}_l],$$

$$\mathbb{B}_s v' = \tilde{f}'[\tilde{P}'_1, \dots, \tilde{P}'_l].$$

Lemma 4.9. *We have*

$$\frac{\tilde{f}}{\tilde{f}'} = S^{l+s} \left(\frac{f}{f'} \right).$$

Proof. As in Lemma 4.8, define the factors g_j, g'_j corresponding to v, v' , respectively. By (4.7) and $(\mathbf{C3})_s$ we have

$$g_l = z^{-1} S^{s+1}(P_1/P_l), \quad (4.9)$$

$$g_{l+j} = z S^{s+1-j}(P_l/P_1) \quad (1 \leq j \leq s), \quad (4.10)$$

and likewise for g'_j .

From (4.6), (4.9) and $(\mathbf{C2})_s$ we obtain

$$S^{l-1}(g'_l) = q^{-l+1} z^{-1} S^{l+s}(P'_1/P'_l) = q^{s+1} S^{l+s}(P_2/P_1) = g_1.$$

Similarly we find

$$S^{j-1}(g_j) = \begin{cases} S^{j+l-2}(g'_{j+l-1}) & (1 \leq j \leq s+1), \\ S^{j-l}(g'_{-l+1}) & (l \leq j \leq l+s). \end{cases}$$

It is clear from $(\mathbf{C1})_s$ and (4.8) that

$$g_j = g'_j \quad (s+2 \leq j \leq l-1).$$

The lemma follows from these relations. $\square$

Lemma 4.10. *Consider a product $\mathbb{B}_s \mathbb{B}_t$ of cancellation blocks $\mathbb{B}_s = C_1 \cdots C_{l+s}$ and $\mathbb{B}_t = D_1 \cdots D_{l+t}$. We assume that either (i) $t = l-2$, or (ii) $l \geq s+t+3$ and $D_{l-s} = \cdots = D_{l-1} = A$. If (v, v') is a pair of vectors satisfying $(\mathbf{C})_t$, then the pair $(\tilde{v}, \tilde{v}') = (\mathbb{B}_t v, \mathbb{B}_t v')$ satisfies $(\mathbf{C})_s$.*

Proof. Set $v = f[P_1, \dots, P_l]$, $v' = f'[P'_1, \dots, P'_l]$. We check $(\mathbf{C})_s$ for $\tilde{v} = \tilde{f}[\tilde{P}_1, \dots, \tilde{P}_l]$ and $\tilde{v}' = \tilde{f}'[\tilde{P}'_1, \dots, \tilde{P}'_l]$, by using Lemma 4.7.

From $(\mathbf{C3})_s$, Lemma 4.7 (ii) and (iii), we have

$$\tilde{P}_1 = zS^{l+t}(P_1), \quad (4.11)$$

$$\tilde{P}_l = q^{l-1}zS^{l+t}(P_{l-t}) = q^{l-1}zS^{l+t}(P_l), \quad (4.12)$$

$$\tilde{P}_j/\tilde{P}_1 = q^{t+1}S^{l+t}(P_2/P_1) \quad (2 \leq j \leq t+2). \quad (4.13)$$

The same relations hold by replacing $P_j, \tilde{P}_j$ with $P'_j, \tilde{P}'_j$. Combining them with $P_1/P_2 = zP'_1/P'_2$ we obtain $\tilde{P}_1/\tilde{P}_2 = z\tilde{P}'_1/\tilde{P}'_2$, which proves $(\mathbf{C2})_s$.

Let us show $(\mathbf{C1})_s$. Note that $D_j = A$ for $2 \leq j \leq t+1$. If $D_j = B$ with $j \geq t+2$, then $\tilde{P}_j = q^{j-1}zS^{l+t}(P_{j-t})$ by Lemma 4.7 (ii), and likewise for $\tilde{P}'_j$. Hence $(\mathbf{C1})_t$ implies $\tilde{P}_j = \tilde{P}'_j$. If $D_i = A$, then we can find some j with $j \geq i+1, t+2$ such that $D_i = D_{i+1} = \cdots = D_{j-1} = A$ and $D_j = B$. Therefore, by Lemma 4.7 (i),

$$\tilde{P}_i = \tilde{P}_j = \tilde{P}'_j = \tilde{P}'_i.$$

Finally, if $t = l-2$, one can show $(\mathbf{C3})_s$ directly since $\mathbb{B}_t = BA^{l-2}B^{l-2}$, and otherwise by using Lemma 4.7 (i) and the assumption $D_{l-s} = \cdots = D_{l-1} = A$. $\square$

4.4. Proof of Proposition 4.2. We now prove Proposition 4.2. For a cancellation pair (ANK, BNK) , we use the symbols g, g', h, h' and Q_i, Q'_i, R_i, R'_i as given in (4.1)–(4.4).

Lemma 4.11. *The vectors $[R_1, \dots, R_l]$ and $[R'_1, \dots, R'_l]$ satisfy $(\mathbf{C})_{s_p}$.*

Proof. For any v and D_i , $D_1 \cdots D_{l-t-1}A^t v$ has the form $f[P_1, \dots, P_l]$ with $P_{l-t} = \cdots = P_l$. Lemma 4.5 implies that Kv_∞ has the same form with $t = s_p + 1$. From this follow $(\mathbf{C2})_s$ and $(\mathbf{C3})_s$. The relation $(\mathbf{C1})_s$ is clear from the definition of A . $\square$

Proof of Proposition 4.2. We set $BN = \mathbb{B}_{s_1} \cdots \mathbb{B}_{s_p}$. Lemma 4.11 shows that $(v, v') = (AKv_\infty, BKv_\infty)$ satisfies $(\mathbf{C})_{s_p}$. By Lemma 4.10, we see inductively that the pair

$$(\tilde{v}, \tilde{v}') = (\mathbb{B}_{s_{i+1}} \cdots \mathbb{B}_{s_p} v, \mathbb{B}_{s_{i+1}} \cdots \mathbb{B}_{s_p} v')$$

satisfies $(\mathbf{C})_{s_i}$ for all i .

Statement (i) is an immediate consequence of Lemma 4.10. Statement (ii) follows from Lemma 4.7 (iii) by taking $\alpha = \sum_{i=1}^p (s_i + 1)$. Finally, from Lemma 4.9 we conclude that $g/g' = S^m(h/h')$ where $m = \sum_{i=1}^p (s_i + l)$. This completes the proof of Proposition 4.2. $\square$

5. BOSONIC FORMULAS

In this section we will prove that the sum of contributions from good monomials is a finite sum of $(l-1)$ -fold series (Corollary 5.7). We parameterize the set of good monomials by a permutation σ and a set of integers $(n_1, \dots, n_{l-1})$ with some restrictions. The vector and scalar part of the corresponding simple vector is explicitly given in terms of these data.

5.1. Parameterization of good monomials. Consider the set $\mathcal{R}$ consisting of $(\sigma, \mathbf{n})$ where $\sigma \in \mathfrak{S}_l$ is such that $\sigma(1) = l$ and $\sigma(l) = 1$, and $\mathbf{n} = (n_1, \dots, n_{l-1}) \in \mathbf{Z}^{l-1}$ satisfying the following conditions:

$$n_i \geq 0 \quad (1 \leq i \leq l-2), \quad (5.1)$$

$$n_{l-1} \geq 2 - \sigma(l-1), \quad (5.2)$$

$$n_i > 0 \quad \text{if} \quad \sigma(i) < \sigma(i+1). \quad (5.3)$$

Proposition 5.1. *There is a one-to-one correspondence between the set $\mathcal{R}$ and the set of good monomials $\mathcal{G}$.*

Proof. We choose the representatives of good monomials of the form $M = C_1 \cdots C_n$ where $C_n = B$. First we construct a mapping $\mathcal{R} \rightarrow \mathcal{G}$. Suppose that $(\sigma, \mathbf{n}) \in \mathcal{R}$. We define for $i = 1, \dots, l-2$

$$\mathbb{E}_i = C_2^{(i)} \cdots C_l^{(i)} \quad (5.4)$$

where

$$C_j^{(i)} = \begin{cases} B & \text{if and only if } j \in \{\sigma(1), \dots, \sigma(i)\}; \\ A & \text{otherwise.} \end{cases} \quad (5.5)$$

We set

$$M = B^{n_{l-1}} \mathbb{E}_{l-2}^{n_{l-2}} \cdots \mathbb{E}_1^{n_1}. \quad (5.6)$$

When $n_{l-1} < 0$ we understand this monomial in the following sense. Set

$$r = \max\{a; 1 \leq a \leq l-2, n_a \neq 0\}. \quad (5.7)$$

Because of (5.3) we have $\sigma(r+1) > \dots > \sigma(l-1)$ and $C_j^{(r)} = A$ if and only if j is equal to one of these $l-r-1$ integers. We have, in particular, that the first (i.e., left) $\sigma(l-1)-2$ elements in $\mathbb{E}_r$ are B . Therefore, by (5.2) we can find M such that $B^{-n_{l-1}} M = \mathbb{E}_r^{n_r} \cdots \mathbb{E}_1^{n_1}$.

In order to show that M given by (5.6) is good it is enough to show that if $C_j^{(i-1)} = B$ then $C_j^{(i)} = B$. This is clear because the only place where $C_j^{(i-1)} \neq C_j^{(i)}$ occurs is at $j = \sigma(i)$, i.e., we have $C_{\sigma(i)}^{(i-1)} = A$ and $C_{\sigma(i)}^{(i)} = B$.

Next, we construct the inverse mapping $\mathcal{G} \rightarrow \mathcal{R}$.

Suppose $M = C_1 \cdots C_n \in \mathcal{G}$. We modify M to $M' = B^{l-1}M$, which is still good. We construct $(\sigma, \mathbf{n})$ for M' by the procedure given below. Then, the one corresponding to M is obtained from $(\sigma, \mathbf{n})$ by changing n_{l-1} to $n_{l-1} - l + 1$.

We write $M' = B^m \mathbb{D}_t \cdots \mathbb{D}_1$ where $m \geq 0$ and $\mathbb{D}_s = D_2^{(s)} \cdots D_l^{(s)}$ and all $\mathbb{D}_s$ contains at least one A . This is, in general, possible because of the modification of M to M' in the first step.

Because of the condition (2.16), we have $D_j^{(s)} = B$ if $D_j^{(s-1)} = B$. For convenience we define $\mathbb{D}_{t+1} = B^{l-1}$. Namely, $D_j^{(t+1)} = B$ for all $2 \leq j \leq l$. We say $D_j^{(s)} = B$ is new if $D_j^{(s-1)} = A$. Besides, all B in $\mathbb{D}_1$ are new by definition. We define $\sigma(i)$ for $1 \leq i \leq l-1$ so that new B 's in $\mathbb{D}_{t+1} \cdots \mathbb{D}_1$, when they are read from right to left, are $D_{\sigma(1)}^{(s_1)}, \dots, D_{\sigma(l-1)}^{(s_{l-1})}$ for some s_i such that $s_1 \leq \dots \leq s_{l-1}$. We set $\sigma(l) = 1$. Note that $\sigma(1) = l$ because $C_n = B$. Thus we obtained an element σ of $\mathfrak{S}_l$.

Now, $\mathbb{E}_i$ and n_i for $1 \leq i \leq l-2$ are uniquely determined by the condition that $\mathbb{D}_t \cdots \mathbb{D}_1$ is of the form $\mathbb{E}_{l-2}^{n_{l-2}} \cdots \mathbb{E}_1^{n_1}$ where (5.5) is satisfied.

We have $M' = B^m \mathbb{E}_{l-2}^{n_{l-2}} \cdots \mathbb{E}_1^{n_1}$. Therefore, we have $M = B^{m-l+1} \mathbb{E}_{l-2}^{n_{l-2}} \cdots \mathbb{E}_1^{n_1}$, and define $n_{l-1} = m - l + 1$.

By the construction the conditions (5.1) and (5.3) are satisfied. Note that the first $\sigma(l-1)-2$ elements of $\mathbb{D}_t \cdots \mathbb{D}_1$ are B . Since $M' = B^{l-1}M$, it implies $\sigma(l-1)-2+m \geq l-1$. Therefore, the condition (5.2) is satisfied.

We have constructed mappings $\mathcal{R} \rightarrow \mathcal{G}$ and $\mathcal{G} \rightarrow \mathcal{R}$. By comparing two constructions we can prove that they are inverse to each other. $\square$

5.2. Actions of good monomials on the vector part. The aim of this section is to describe the actions of good monomials M given by (5.6) where we consider A and B as operators acting on l -component vectors $[P_1, \dots, P_l]$ by

$$\begin{aligned} A : [P_1, \dots, P_l] &\mapsto [S(P_1), S(P_1), S(P_2), \dots, S(P_{l-1})], \\ B : [P_1, \dots, P_l] &\mapsto [zS(P_l), S(P_1), \dots, S(P_{l-1})]. \end{aligned}$$

Namely, these actions are obtained from that of A and B forgetting the scalar part. We call it the action on the vector part and denote by the same letters A, B .

First we describe the action of B^m on the vector part. Suppose $[P_1, \dots, P_l]$ is given. Define $P_i^{(n)}$ for all $i, n \in \mathbf{Z}$ by

$$P_{j+ml} = q^{-\frac{1}{2}m(m-1)l-(j-1)m} z^{-m} P_j, \quad (5.8)$$

$$P_i^{(n)} = S^n(P_{i-n}). \quad (5.9)$$

Lemma 5.2. *The formula (5.8) is consistent. The action of B^n ($n \in \mathbf{Z}$) on the vector part is given by*

$$B^n[P_1, \dots, P_l] = [P_1^{(n)}, \dots, P_l^{(n)}]. \quad (5.10)$$

Proof. We define P_i for arbitrary $i = j + ml \in \mathbf{Z}$ by (5.8) where $1 \leq j \leq l$ and $m \in \mathbf{Z}$. The consistency of (5.8) can be checked by induction on j for all j , reducing the case $j > l$ to $j - l$, and $j < 0$ to $j + l$. We then define $P_i^{(n)}$ for all $i, n \in \mathbf{Z}$ by (5.9). The validity of (5.10) can be checked by induction on n , reducing $n > 0$ to $n - 1$, and $n < 0$ to $n + 1$. $\square$

The following formula follows from $S \circ z = qz \circ S$.

$$P_i^{(n)} = q^{\frac{1}{2}m(m+1)l+(1-i)m} z^{-m} S^n(P_j) \quad \text{where } i - n = j + ml. \quad (5.11)$$

Next we describe the action of $\mathbb{E}_i^{n_i}$ on the vector part. For this purpose we prepare a few notations. We consider a subset $I \subset [2, l]$ such that $l \in I$. Let r be the cardinality of I . We identify the subset I with a mapping $I : \mathbf{Z} \rightarrow [1, l]$ by

$$\begin{aligned} I(0) &= 1, \\ I &= \{I(1), \dots, I(r)\} \quad \text{where } I(1) < \dots < I(r), \\ I(i + r + 1) &= I(i). \end{aligned}$$

Note that $I(r) = l$. If $r = l - 1$, we have $I(i) = i + 1$ for $0 \leq i \leq l - 1$.

We also define $I^* : [1, l] \rightarrow [0, r]$ by

$$I^*(j) = \max\{i; 0 \leq i \leq r, I(i) \leq j\}. \quad (5.12)$$

Note that $I^* \circ I(i) = i$ ($0 \leq i \leq r$), $I^*(1) = 0$ and $I^*(l) = r$. If $r = l - 1$, we have $I^*(j) = j - 1$.

Let I be as above. We define an operator $B[I] = C_2 \cdots C_l$ where $C_i = A$ or B and $C_i = B$ if and only if $i \in I$. As we will see in the following proposition, the action of $B[I]^n$ on the vector part $[P_1, \dots, P_l]$ depends only on P_i ($i \in I$), and the resulting vector part $[P'_1, \dots, P'_l]$ has the property $P'_j = P'_{I \circ I^*(j)}$. Namely, the action of $B[I]^n$ on the vector part is actually understood as acting on the $r + 1$ component vector $P_{I(i)}$ ($0 \leq i \leq r$). This is the reason for introducing the mappings I and I^* .

Lemma 5.3. *Let I and r be as above and $[P'_1, \dots, P'_l] = B[I]^n[P_1, \dots, P_n]$ ($n \in \mathbf{Z}_{\geq 0}$). We have*

$$P'_j = q^{\omega_r(I, n, i)} z^{\rho_r(n, i)} S^{(l-1)n}(P_{I(i+n)}) \quad \text{where } i = I^*(j), \quad (5.13)$$

$$\begin{aligned} \omega_r(I, n, i) &= \sum_{s=1}^n (I(i+s) - 2) + \frac{1}{2}n(n-1)(l-1) \\ &\quad + m - (l-1)\{(r-i)m + \frac{1}{2}m(m-1)(r+1)\}, \end{aligned} \quad (5.14)$$

$$\rho_r(n, i) = n - m, \quad (5.15)$$

where $m = m(n, i)$ denotes the integer part of $(i+n)/(r+1)$.

Proof. For $n = 1$ we have

$$\begin{aligned} P'_{I(i)} &= \cdots = P'_{I(i+1)-1} = q^{I(i+1)-2} z S^{l-1}(P_{I(i+1)}) \quad (0 \leq i \leq r-1), \\ P'_{I(r)} &= P'_l = S^{l-1}(P_1). \end{aligned}$$

Set

$$S(i) = \begin{cases} q^{I(i)-2} z S^{l-1} & \text{if } i \not\equiv 0 \pmod{r+1}; \\ S^{l-1} & \text{otherwise.} \end{cases}$$

The general case reads as

$$P'_{I(i)} = S(i+1) \circ \cdots \circ S(i+n)(P_{I(i+n)}).$$

The result (5.13) is obtained by using the commutation relations $S \circ z = qz \circ S$. The number of times that $i+s \equiv 0 \pmod{r+1}$ occurs for $1 \leq s \leq n$ is given by $m(n, i)$. For such s the counting of the powers of q and z differs from $i+s \not\equiv 0 \pmod{r+1}$. Taking this difference into account we obtain (5.13). $\square$

Proposition 5.4. *Let M be a good monomial and $[P_1, \dots, P_l]$ the vector part of $M[1, \dots, 1]$. Let $(\sigma, \mathbf{n})$ correspond to M as in Proposition 5.1. Fix $1 \leq i \leq l$ and define j and m by $i - n_{l-1} = j + ml$ and $1 \leq j \leq l$. Define I_a, i_a ($1 \leq a \leq l-2$) by*

$$I_a = \{\sigma(1), \dots, \sigma(a)\} \quad i_a = \begin{cases} I_{l-2}^*(j) & \text{if } a = l-2; \\ I_a^* \circ I_{a+1}(i_{a+1} + n_{a+1}) & \text{if } 1 \leq a \leq l-3, \end{cases}$$

(see (5.12)). Then, we have

$$P_i = q^{\omega(\sigma, \mathbf{n})_i} z^{\rho(\sigma, \mathbf{n})_i}, \quad (5.16)$$

$$\begin{aligned} \omega(\sigma, \mathbf{n})_i &= \frac{1}{2}m(m+1)l + (1-i)m + \sum_{a=1}^{l-2} \omega_a(I_a, n_a, i_a) \\ &\quad + n_{l-1} \sum_{a=1}^{l-2} \rho_a(n_a, i_a) + (l-1) \sum_{1 \leq a < b \leq l-2} n_b \rho_a(n_a, i_a), \end{aligned} \quad (5.17)$$

$$\rho(\sigma, \mathbf{n})_i = -m + \sum_{a=1}^{l-2} \rho_a(n_a, i_a). \quad (5.18)$$

Proof. This proposition follows from Lemma 5.2 and Lemma 5.3. $\square$

Consider $\omega(\sigma, \mathbf{n})_i$ and $\rho(\sigma, \mathbf{n})_i$ as polynomials of $\mathbf{n} = (n_1, \dots, n_{l-1})$. They depend on σ and i . However, the quadratic part $\omega^{(2)}(\mathbf{n})$ in $\omega(\sigma, \mathbf{n})_i$ and the linear part $\rho^{(1)}(\mathbf{n})$ in $\rho(\sigma, \mathbf{n})_i$ are independent:

$$\begin{aligned} \omega^{(2)}(\mathbf{n}) &= \frac{l-1}{2} \sum_{\substack{1 \leq a, b \leq l-2 \\ c = \min(a, b)}} n_a n_b \frac{c}{c+1} + n_{l-1} \sum_{1 \leq a \leq l-2} n_a \frac{a}{a+1} + \frac{n_{l-1}^2}{2l}, \\ \rho^{(1)}(\mathbf{n}) &= \sum_{1 \leq a \leq l-2} n_a \frac{a}{a+1} + \frac{n_{l-1}}{l}. \end{aligned}$$

5.3. Coloring blocks and scalar factors. In this section, we determine the scalar part of the monomial Mv_∞ . It is of the form

$$\frac{(-1)^\alpha q^\beta z^\gamma}{(q)_{t_2} \cdots (q)_{t_l} (q^{\sum_{c=2}^l t_c z})_\infty}$$

We will determine α, β, γ and t_a ($2 \leq l$) for each $M = C_1 \cdots C_n$.

A sub-monomial of the form $C_j \cdots C_{j+r-1}$ ($1 \leq r \leq l-1, 1 \leq j \leq n+1-r$) is called a block if $j = n, r = 1$ or the following conditions are satisfied:

$$C_i = \begin{cases} B & \text{for } i = j, j+r; \\ A & \text{for } j+1 \leq i \leq j+r-1. \end{cases}$$

Namely, a block is a maximal (i.e., $C_{j+r} = B$) submonomial of the form $BA \cdots A$. The restriction $r \leq l-1$ is a consequence of the assumption that M is good and $C_n = B$.

Set $\mathbf{B} = \{j; 1 \leq j \leq n, C_j = B\}$. Obviously, the blocks in M are labeled by $\mathbf{B}$. We define the length and color of each block by mappings $r : \mathbf{B} \rightarrow \{1, \dots, l-1\}$ and $c : \mathbf{B} \rightarrow \{1, \dots, l\}$. The length $r(j)$ of the block $C_j \cdots C_{j+r-1}$ is r . The color of each block is defined inductively as follows. We set $c(n) = 1$. Suppose we have defined $c(j)$ for all $j \in \mathbf{B}$ such that $i < j$. Set $i_1 = i + r(i) \in \mathbf{B}$. If $i_1 + l - 1 > n$ or $C_{i_1+l-1} = A$, we set $c(i) = \max\{c(j); j \in \mathbf{B}, i < j\} + 1$. Otherwise, we have $i_1 + l - 1 \in \mathbf{B}$; we set $c(i) = c(i_1 + l - 1)$.

It is convenient to show the color c of B in M by B_c . For example, if $l = 4$, the coloring of $M = B^5(BAB)^4(AAB)^3$ is

$$B_3 B_1 B_4 B_2 B_3 B_1 A B_2 B_3 A B_1 B_2 A B_3 B_1 A B_2 A A B_1 A A B_2 A A B_1.$$

Fix a monomial M . Define $\lambda : \mathbf{B} \rightarrow \mathbf{Z}_{\geq 0}$ as follows.

$$\lambda_i = \begin{cases} \sum_{\substack{j \in \mathbf{B} \\ i < j, c(j) \neq 1}} r(j) & \text{if } c(i) = 1; \\ \sum_{\substack{j \in \mathbf{B} \\ i \leq j, c(j) = c(i)}} r(j) & \text{if } c(i) \neq 1. \end{cases} \quad (5.19)$$

Namely, if $c(i) = 1$, λ_i is the total length of the blocks of color not equal to 1 that are located to the right of the block i . If $c(i) \neq 1$, λ_i is the total length of the blocks of color $c(i)$ that are located to the right of the block i including itself.

Lemma 5.5. *If $C_i = C_j = B_c$ where $i < j$, then we have $i + l \leq j$.*

Proof. We prove the statement by induction on i starting from $i = n$ where the statement is valid because the “if” statement is false. Suppose $C_i = C_j = B_c$ where $i < j$ and $i + l > j$. Let i' be the smallest integer such that $i < i'$ and $C_{i'} = B$. Since the color c is attained at j before i , by the way of coloring we have $i' + l - 1 \leq n$ and $C_{i'+l-1} = B_c$. Note that $i' \leq j < i + l \leq i' + l - 1$, and therefore we have $0 < i' + l - 1 - j \leq l - 1$ and $C_j = C_{i'+l-1} = B_c$, which contradicts to the induction hypothesis. $\square$

The following proposition determines the scalar factors.

Proposition 5.6. *Let M be a good monomial, $M = C_1 \cdots C_n$ and $C_n = B$. Set $M_i = C_i \cdots C_n$ and $M_i v_\infty = f_i[P_1^{(i)}, \dots, P_n^{(i)}]$. We set formally $M_{n+1} v_\infty = v_\infty = f_{n+1}[1, \dots, 1]$, where $f_{n+1} = \frac{1}{(z)_\infty}$. For $1 \leq i \leq n$, define $i_b \in \mathbf{B}$ such that $i_b \leq i \leq i_b + r(i_b) - 1$. Then, we have*

$$h_i = \frac{f_i}{S(f_{i+1})} = \begin{cases} 1/(1 - q^{\lambda_{i_b}} z) & \text{if } C_i = A \text{ and } c(i_b) = 1; \\ 1/(1 - q^{-\lambda_{i_b}} z^{-1}) & \text{if } C_i = B \text{ and } c(i_b) = 1; \\ 1/(1 - q^{-\lambda_{i_b} + i - i_b}) & \text{if } C_i = A \text{ and } c(i_b) \neq 1; \\ 1/(1 - q^{\lambda_{i_b} - i + i_b}) & \text{if } C_i = B \text{ and } c(i_b) \neq 1. \end{cases} \quad (5.20)$$

Proof. We prove the statement by induction on i_b starting from $i_b = n$. Our induction hypotheses are the following.

(i) The vector part of $M_{i_b+r(i_b)+1} v_\infty$ is of the form $[q^{\alpha_1} : \dots : q^{\alpha_l}]$ if $c(i_b) = 1$, otherwise it is $[q^{\alpha_1} z : \dots : q^{\alpha_l}]$;

In the former case, all components are of the form q^{α_j} , and in the latter there exists $1 \leq j_0 < l$ such that $q^{\alpha_j} z$ if $j \leq j_0$ and q^{α_j} if $j > j_0$.

$$(ii) \alpha_l - \alpha_1 = \begin{cases} \lambda_i & \text{if } c(i_b) = 1; \\ -\lambda_i & \text{otherwise.} \end{cases}$$

In the course of induction, we will also prove (5.20).

If $i_b = n$, then $i = n$ and $\lambda_n = 0$. We have $c(n) = 1$ and $h_n = 1 - z^{-1}$ by (2.13), (2.14) and (2.8). Note also that the vector part of $M_{n+1} v_\infty$ is $[1, \dots, 1]$.

In the below we use the projective notation $[Q_1 : \dots : Q_l]$ for the vector part $[P_1, \dots, P_l]$. Namely, $Q_i = Q P_i$ where Q is independent of i . We use the symbols $\uparrow$, $\downarrow$ and $|$ to indicate the submonomial sitting in the right of these symbols. For example, if we write $C_1 \cdots C_i \uparrow C_{i+1} \cdots C_n$, the vector part at $\uparrow$ means that of $C_{i+1} \cdots C_n v_\infty$.

Consider i such that $c(i_b) = 1$. The monomial M_{i_b} is of the form

$$\underbrace{B_1 A \cdots A}_s \uparrow \underbrace{B \cdots B A \cdots A}_{l-1} B_1 \downarrow C_{i+s+l} \cdots C_n.$$

Here $A \cdots A$ are all A , but $B \cdots B$ can be a mixture. We have the restriction $s \leq r$ because M is good. Note that C_i belongs to $\underbrace{B_1 A \cdots A}_s$.

By the induction hypothesis the vector part is of the form $[q^{\alpha_1} : \dots : q^{\alpha_l}]$ at $\downarrow$. Then, by the following B_1 , the vector part changes to $[q^{\alpha_l} z : q^{\alpha_1} : \dots]$, and after the next $r - 1$ of A , it further changes to

$$\underbrace{[q^{\alpha_l+r-1} z : \dots : q^{\alpha_l+r-1} z : q^{\alpha_1} : \dots]}_r.$$

At the point $\uparrow$ it is of the form

$$[q^{\alpha_1} : \dots : \underbrace{q^{\alpha_l+l-1} : \dots : q^{\alpha_l+l-1}}_r].$$

The assertion (5.20) and the statement (i) immediately follow from this.

Note that the difference $\alpha_l - \alpha_1$ have increased by $l - 1$ from $\downarrow$ to $\uparrow$. By Lemma 5.5 there is only one B_1 between $\uparrow$ and $\downarrow$. Therefore, the total length of blocks of color not 1 between $\uparrow$ and $\downarrow$ is equal to $l - 1$. Therefore, the statement (ii) is also proved.

Next, we consider the case $c(i_b) = c$ where $c \neq 1$. There are two cases: (a) $C_j \neq B_c$ for all $j > i_b$, and (b) there exists $C_j = B_c$ for some $j > i_b$. In the latter case, we have $i_b < j - l + 1$, $C_{j-l+1} = B$ and $C_{i'} = A$ for $i_b < i' < j - l + 1$.

Case (a): the monomial M_i is of the form

$$\underbrace{B_c \uparrow A \cdots A}_{s} \mid \overbrace{B \cdots \downarrow}^t \overbrace{B \cdots B A \cdots A}^{l-1} \underbrace{B \cdots}_{r} .$$

Here $A \cdots A$ are all A , but other dots can be A or B . We have $s \leq r$ because M is good. C_i belongs to the block $\underbrace{B_c \uparrow A \cdots A}_{s-t}$.

At the point $\downarrow$ the vector part is of the form $[\cdots : \underbrace{1 : \cdots : 1}_r]$. It changes to $[z : \cdots : \underbrace{1 : \cdots : 1}_{r-t}]$ at $\mid$, and then to $[q^{s-t-1}z : \cdots : \underbrace{1 : \cdots : 1}_{r-s}]$ at $\uparrow$. The assertion (5.20) and statements (i) and (ii) follow from these observations.

Case (b): the monomial M_i is of the form

$$\underbrace{B_c \uparrow A \cdots A}_{s} \mid \overbrace{B \cdots B A \cdots A}^{l-1} \underbrace{B_c \downarrow \cdots}_{r} ,$$

where $s \leq r$ and there are no B_c between the two B_c in this diagram. By the induction hypothesis, the vector part at the point $\downarrow$ is of the form $[q^{\alpha_1}z : \cdots : q^{\alpha_l}]$. It changes to $[q^{\alpha_1+1}z : \cdots : \underbrace{q^{\alpha_l} : \cdots : q^{\alpha_l}}_r]$ at $\mid$, and then to $[\underbrace{q^{\alpha_1+s}z : \cdots : q^{\alpha_1+s}z}_s : \cdots : \underbrace{q^{\alpha_l} : \cdots : q^{\alpha_l}}_{r-s+1}]$ at $\uparrow$. The assertion (5.20) and the statements (i) and (ii) follow from these observations. $\square$

For a given good monomial $M = C_1 \cdots C_n$ with $C_n = B$ we define t_c and m_c ($1 \leq c \leq l$): the total length of color c blocks is denoted by t_c , and the number of color c blocks is denoted by m_c . We also set

$$\mu_c = \sum_{i: C_i = B_c} (n - i + 1).$$

Corollary 5.7. *The scalar part f of $Mv_\infty = C_1 \cdots C_n v_\infty = f[P_1, \dots, P_l]$ is given by the formula*

$$f = \frac{(-1)^\alpha q^{\bar{\beta}} z^{m_1}}{(q)_{t_2} \cdots (q)_{t_l} (q^{\sum_{c=2}^l t_c z})_\infty}, \quad (5.21)$$

$$\alpha = n - t_1 + \sum_{c=1}^l m_c, \quad (5.22)$$

$$\bar{\beta} = nm_1 + \frac{1}{2} \sum_{c=2}^l t_c(t_c + 1) - \sum_{c=1}^l \mu_c. \quad (5.23)$$

As in the case of vector part, the quadratic part of $\bar{\beta}$ and the linear parts of $m_1, t_2, \dots, t_l$ are independent of σ . This follows from the following formula for the linear parts. Below $x \sim y$ denotes an equality modulo constant term in $n_1, \dots, n_{l-2}$.

$$\begin{aligned} n &= (l-1) \sum_{a=1}^{l-2} n_a + n_{l-1}, \\ m_c &\sim \sum_{a=c-1}^{l-2} \frac{a}{a+1} n_a + \frac{n_{l-1}}{l}, \\ t_c &\sim (l-1) \sum_{a=c-1}^{l-2} \frac{n_a}{a+1} + \frac{n_{l-1}}{l}. \end{aligned}$$

Acknowledgments. This work is partially supported by the Grant-in-Aid for Scientific Research (B) no.12440039 and (A1) no.13304010, Japan Society for the Promotion of Science. The work of SL is in part supported by grant RFBR-01-01-00546

REFERENCES

- [B] R. Baxter, *Exactly solved models in statistical mechanics*, Academic Press, London, 1982.
- [FJM] B. Feigin, M. Jimbo and T. Miwa, *Vertex operator algebra arising from the minimal series $M(3, p)$ and monomial basis*, RIMS-**1310**, math.QA/0012193.
- [BMS] A. Berkovich, B. McCoy and A. Schilling, *Rogers-Schur-Ramanujan type identities for the $M(p, p')$ minimal models of conformal field theory*, Commun. Math. Phys. **191** (1998), 325–395.
- [FS] B. Feigin and A. Stoyanovsky, *Quasi-particles models for the representations of Lie algebras and geometry of flag manifold*, hep-th/9308079, RIMS 942; *Functional models for the representations of current algebras and the semi-infinite Schubert cells*, Funct. Anal. Appl. **28** (1994), 55–72.
- [FL] B. Feigin and S. Loktev, *On finitization of the Gordon identities*, math.QA/0006221.
- [JMM] M. Jimbo, T. Miwa and E. Mukhin, *Bosonic formula for $\widehat{sl}_2$ coinvariants*, math.QA/0107053.
- [PKh] A. Pukhlikov, A. Khovanskii, *Finitely additive measures of virtual polytopes*, St. Petersburg Math J. **4** (1993), 337–356.

BF: LANDAU INSTITUTE FOR THEORETICAL PHYSICS, CHERNOGOLOVKA, 142432, RUSSIA
 E-mail address: feigin@feigin.mccme.ru

MJ: GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, TOKYO 153-8914, JAPAN

E-mail address: jimbomic@ms.u-tokyo.ac.jp

SL: INSTITUTE FOR THEORETICAL AND EXPERIMENTAL PHYSICS AND INDEPENDENT UNIVERSITY OF MOSCOW

E-mail address: loktev@mccme.ru

TM: DIVISION OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, KYOTO UNIVERSITY, KYOTO 606-8502 JAPAN

E-mail address: tetsuji@kusm.kyoto-u.ac.jp

EM: DEPT. OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720

E-mail address: mukhin@math.berkeley.edu